

INPFC
DOCUMENT
Sec. No. <del>    </del>
Rev. No. 2239

BIOCHEMICAL GENETIC VARIATION IN WALLEYE POLLOCK, THERAGRA CHALCOGRAMMA:

II. ANALYSIS OF PELAGIC COLLECTIONS OF 1978 FROM THE CENTRAL AND SOUTHERN  
BERING SEA (ABSTRACT OF REPORT IN PROGRESS)

by

Fred Utter, George Milner, and David Teel

Submitted to the

INTERNATIONAL NORTH PACIFIC FISHERIES COMMISSION

by the U.S. NATIONAL SECTION

National Marine Fisheries Service  
Northwest and Alaska Fisheries Center  
2725 Montlake Boulevard East  
Seattle, Washington 98112

October 1979

BIOCHEMICAL GENETIC VARIATION IN WALLEYE POLLOCK, THERAGRA CHALCOGRAMMA:

II. ANALYSIS OF PELAGIC COLLECTIONS OF 1978 FROM THE CENTRAL AND SOUTHERN  
BERING SEA (ABSTRACT OF REPORT IN PROGRESS)

by Fred Utter\*, George Milner\*, and David Teel\*

This report summarizes biochemical genetic data from 324 individuals from the central and southern Bering Sea (Fig. 1), and 27 individuals from the North Pacific Ocean near Japan. The frozen storage of muscle, heart, and liver tissues from these fish for over one year resulted in significant deterioration of much enzyme activity, and precluded a definitive comparison of allelic frequencies from these collections with similar data from collections of pollock from other areas. Nevertheless, the data are valuable as a source of (1) previously undescribed genetic variation, (2) some comparative data with other studies, and (3) suggestions of structuring that require confirmation by subsequent studies.

The only useful data were from extracts of skeletal muscle, although all three tissues were extensively screened. These data included 14 monomorphic loci [fluorescent acid phosphatase (ACP), alphaslycerophosphate dehydrogenase (AGP), esterase (EST)-3, general protein (GP) - three loci, glyoxilase (GLO), isocitrate dehydrogenase (IDH), malic enzyme (ME)-1, malic dehydrogenase (MDH)-3, peptidase (PEP)-2, PEP-6, phosphoglucomutase (PGM), phosphoglucose isomerase (PGI)-2, and 11 polymorphic loci (Tables 1 and 2) adeonsine deaminase (ADA)-1, EST-1, lactate dehydrogenase (LDH)-1, LDH-2, LDH-3, MDH-1, MDH-2, ME-2, PEP-4, PEP-5, PGI-3]. Nineteen apparent or putative loci were also tentatively identified where electrophoretic expression was either indistinct, unreliable or weak [aconitase (ACON),

---

\* National Marine Fisheries Service, Northwest and Alaska Fisheries Center, Seattle, Washington 98112.

ADA-2, aldolase (ALD), EST-2, fumarase (FUM), beta galactosidase (b-GAL), glyceraldehyde-3-phosphate dehydrogenase (GAP), guanine deaminase (GDA), glutamic dyhydrogenase (GLUD), glutamic-pyruvic transaminase (GPT), glutathione reductase (GR), leucine aminopeptidase (LAP), nucleoside phosphorylase (NP), 6-phosphogluconate dehydrogenase (6-PGD), phosphomannose isomerase (PMI), PEP-1, PEP-3, tetraoxolium oxidase (TO), xanthine dehydrogenase (XDH)]. Both PMI and 6-PGD in this latter category were obviously highly polymorphic and promise to be useful for population studies when analyzed from fresher materials. Iwata (1975) has reported major differences in the frequency of TO alleles between pollock populations of the eastern and western North Pacific Ocean.

Comparisons of allelic frequencies from five of the polymorphic loci (ME-1, ME-2, MDH-1, LDH-1, and LDH-2) could be made with frequencies reported by Grant and Utter (JFRBC, in press) of collections from the northeastern Pacific Ocean and the southeastern Bering Sea. Allelic frequencies at each of these loci were generally within the ranges of values reported by Grant and Utter (in press), and did not define any new population units.

Allelic frequencies at a previously unreported locus (LDH-3) differed markedly among the collections. The average allelic frequencies of most of the Bering Sea samples differed by a factor of about .50 from the average allelic frequency of the most southern Bering Sea collection (52°20'N, 178°16'E) and the Japanese sample. The preliminary nature of these data precludes further statements concerning this variation. More rigorous testing is required to confirm the genetic nature of this variation, and if confirmed, more sampling is required to determine its oceanic distribution.

It is recommended (1) that future studies include the ability to analyze the samples within a few weeks of collection because of the particularly labile nature of gadid proteins, and (2) that collection points be extended to the northwest Pacific Ocean south of  $52^{\circ}\text{N}$ .

## Polymorphic Loci

Collection	N	LDH-1		LDH-2			LDH-3		EST-1		ADA-1		
		A	B	A	B	C	A	B	A	B	A	B	C
Bering Sea Haul 1	48	1.0	0	1.0	0	0	.86	.14	1.0	0	1.0	0	0
5	48	.99	.01	.99	.01	0	.65	.35	1.0	0	.98	.01	.01
13	48	1.0	0	.99	0	.01	.66	.34	1.0	0	1.0	0	0
33	21	1.0	0	.96	0	.04	.95	.05	1.0	0	1.0	0	0
36	47	1.0	0	.99	.01	0	.69	.31	.99	.01	.99	.01	0
40	23	1.0	0	1.0	0	0	.14	.86	1.0	0	1.0	0	0
67	48	1.0	0	1.0	0	0	.80	.20	1.0	0	1.0	0	0
78	27	1.0	0	1.0	0	0	.98	.02	1.0	0	1.0	0	0
Japanese	27	1.0	0	1.0	0	0	.32	.68	1.0	0	1.0	0	0

	N	MDH-1			MDH-2			ME-2		PEP-4			PEP-5			PGI-3	
		A	B	C	A	B	C	A	B	A	B	C	A	B	C	A	B
Haul 1	48	.99	.01	0	1.0	0	0	.47	.53	1.0	0	0	.94	.06	0	1.0	0
5	48	1.0	0	0	.99	.01	0	.41	.59	.87	.13	0	.96	.04	0	1.0	0
13	48	1.0	0	0	.99	0	.01	.42	.58	.98	0	.02	.99	.01	0	.99	.01
33	21	.95	.05	0	1.0	0	0	--	-	-	-	-	-	-	-	.98	.02
36	47	.99	.01	0	1.0	0	0	.46	.54	.97	.03	0	.98	.02	0	1.0	0
40	23	1.0	0	0	1.0	0	0	.38	.62	1.0	0	0	.95	.05	0	.98	.02
67	48	1.0	0	0	1.0	0	0	.60	.40	.98	.02	0	.86	.13	.01	1.0	0
78	27	.96	.04	0	.98	.02	0	.48	.52	1.0	0	0	-	-	-	1.0	0
Japanese	27	.96	.04	0	1.0	0	0	.43	.57	.98	0	.02	.91	.02	.07	1.0	0

Table 1.--Allelic frequencies of polymorphic systems. Allelic form designated "A" is most frequent for each system and assigned a relative mobility of 100; see Table 2 for relative mobilities of alternate allelic forms.

<u>Locus</u>	<u>Alternate Allele</u>	
	<u>B</u>	<u>C</u>
MDH-1	0.15	0.24
MDH-2	0.73	1.58
ME-2	1.10	
PEP-4	1.27	0.91
PEP-5	1.10	0.89
LDH-3	0.33	
LDH-1	0.63	
LDH-2	0.91	0.67
PGI-3	0.93	
EST-1	1.10	
ADA-1	0.62	1.14

Table 2.--Relative mobilities of alternate allelic forms.




Figure 1. -- Locations of Bering Sea samples.