

NPAFC
Doc. 1508
Rev. _____

Southeast Alaska Coastal Monitoring (SECM) Survey Plan for 2014

by

Joseph A. Orsi, Emily A. Fergusson, Edward V. Farley Jr., and Ronald A. Heintz

Auke Bay Laboratories
Alaska Fisheries Science Center, NMFS,
NOAA Fisheries, U.S. Department of Commerce,
Ted Stevens Marine Research Institute
17109 Point Lena Loop Road
Juneau, AK 99801 USA

Submitted to the

NORTH PACIFIC ANADROMOUS FISH COMMISSION

by

the United States of America

March 2014

THIS PAPER MAY BE CITED IN THE FOLLOWING MANNER:

J. A. Orsi, E. A. Fergusson, E. V. Farley, Jr., and R. A. Heintz. 2014. Southeast Alaska Coastal Monitoring (SECM) Survey Plan for 2014. NPAFC Doc. 1508. 17 pp. (Available at <http://www.npafc.org>).

Southeast Alaska Coastal Monitoring (SECM) Survey Plan for 2014

Keywords: juvenile salmon, ocean monitoring, ecology, Southeast Alaska

ABSTRACT

This survey plan details the proposed sampling for the Southeast Coastal Monitoring (SECM) project in May, June, July, and August of 2014. The SECM project is supported by the Alaska Fisheries Science Center (AFSC), Auke Bay Laboratories (ABL), with an objective to study the habitat use and early marine ecology of juvenile (age-0) Pacific salmon (*Oncorhynchus* spp.) and associated epipelagic ichthyofauna in Southeast Alaska and in the Gulf of Alaska ecosystem. The SECM surveys have been continuous since 1997, and have provided long-term biological and oceanographic data sets associated with all five species of both wild and hatchery salmon during a period of climate change.

SECM sampling planned for 2014

The Southeast Coastal Monitoring (SECM) project was initiated in 1997 by the Alaska Fisheries Science Center (AFSC), Auke Bay Laboratories (ABL), to study the habitat use and early marine ecology of juvenile (age-0) Pacific salmon (*Oncorhynchus* spp.) and associated epipelagic ichthyofauna in Southeast Alaska. SECM surveys are conducted off of government and chartered research vessels in the vicinities of Icy Strait and Icy Point (58° N, 135°W; 57° N, 134° W) to sample fish, zooplankton, nutrients/chlorophyll, and physical water properties using a surface trawl, plankton nets, and an oceanographic profiler. This monitoring has permitted compilation of long-term biological and oceanographic data sets for a region producing all five

species of both wild and hatchery salmon during a period of climate change. In 2014, SECM research surveys are scheduled to continue at 13 core stations during four monthly intervals from late May to late August (Figure 1, Tables 1 and 2). This 2014 SECM survey plan reflects accomplishments since the inception of the project, and outlines how this additional survey year will extend the biophysical time series to 18 years.

SECM researchers have collaborated with many agencies, institutions, and individuals over the project lifespan and have contributed numerous reports and publications to the scientific community (http://www.afsc.noaa.gov/ABL/MSI/msi_secm.htm; see also selected publications section). Annual presentations on pink salmon forecasting have been given to the Purse Seine Task Force, (http://www.afsc.noaa.gov/ABL/MSI/msi_sae_psf.htm) since 2005, and both oral and poster presentations on topics in salmon ecology are made at professional meetings and seminars, such as the Pink and Chum Salmon Workshop, Salmon Ocean Ecology Meeting, American Fisheries Society Alaska Chapter and national meetings, Alaska Hatchery Managers Meeting, North Pacific Anadromous Fish Commission (NPAFC) Meeting, Alaska Marine Science Symposium, State of the Salmon international conference, and at local schools and universities. A complete summary of SECM data for the prior year is reported annually to NPAFC in its document series. These contributions are an important service to the fisheries community and continue to provide data that will improve understanding of the role of salmon in regional and basin-scale ecosystems. The increasing value of these time series has most recently become apparent by providing information on juvenile Chinook salmon ecology that can be applied to improve understanding of stocks in western Alaska, where the causes for disastrous run failures are unknown.

Historical SECM sampling 1997–2013

Historically, some aspects of the SECM project have varied to accommodate new research objectives or process studies designed to examine specific questions. From 1997 to 2000, SECM research was directed at sampling juvenile salmon and their associated biophysical parameters in inshore, strait, and coastal habitats along a primary seaward migration corridor in the northern region of Southeast Alaska. Up to 24 stations spanning 250 km were sampled five times annually, from May to early October. These habitats extended geographically from inshore localities near large glacial rivers to 65 km offshore in the Gulf of Alaska. Fish were sampled during the daytime with a NORDIC 264 surface rope trawl from the National Oceanic and Atmospheric Administration (NOAA) Ship *John N. Cobb*. The biophysical data collected included vertical profile data on water temperature and salinity, surface nutrients and chlorophyll, zooplankton from vertical 20-m hauls and double oblique hauls deployed to 200 m depth, and onboard stomach analysis of potential predators of juvenile salmon. A laboratory process study of juvenile sablefish predation impact on juvenile salmon was initiated in response to unusual field observations in 1999. After four years of sampling, the inshore sampling stations and the Cross Sound transect were eliminated because juvenile salmon trawl catches were consistently low in these habitats during the sampling months.

In 2001–2004, SECM researchers continued biophysical sampling at 13 core stations and directed more research effort into process studies. Two such studies initiated in 2001 included diel feeding periodicity and prey fields of juvenile pink, chum, and coho salmon, and onboard gastric evacuation rate experiments for juvenile pink and chum salmon. These process studies were designed to increase our understanding of trophic linkages and provide more specific input parameters for use in bioenergetics models to evaluate coastal marine carrying capacity and

salmon habitat quality (growth potential). Beginning in 2002, the late September sampling was eliminated because juvenile salmon abundances had been consistently low for this monthly interval, and consequently, sampling time intervals were extended in earlier months to maximize the opportunities for obtaining data at offshore stations and to replicate trawling at the core stations. In 2003, sampling frequency at the 13 core stations was increased from four to six time intervals between mid-May and late August, to gain better temporal resolution of biophysical factors related to salmon growth and abundance. Two process studies were conducted in this year, one on sea lice infestation of juvenile salmon at sea and another on energetics of starved juvenile chum salmon brought into the laboratory from trawl catches during field surveys. Additionally, in 2003 and 2004, concurrent sampling with a second, smaller mesh, trawl was conducted in two time intervals to examine spatial distribution and to compare size-selectivity of two trawl types for juvenile salmon. Also in 2004, the SECM trawl sampling effort returned to one of the inshore habitats to collaborate with a Taku Inlet study on hatchery- and wild-stock interactions of juvenile chum salmon.

From 2005 to 2009, SECM research expanded to include a second sampling region in southern Southeast Alaska. The established sampling scheme of 13 core stations was maintained in the northern region from late May to late August, and eight stations in Clarence Strait were also sampled in late June and late July. The 2008 scheduled sampling in Clarence Strait was not accomplished because of a fatal vessel breakdown of the *John N. Cobb*. In 2009, sampling was resumed in the southern region; however, vessel charter contract problems only allowed sampling to occur in July in this region. Future efforts are not planned for the southern region, but the four years of comparative sampling of juvenile salmon and associated biophysical parameters will be used to broaden basic understanding of the trophic relationships and

ecological interactions of wild and hatchery juvenile salmon and associated fishes in different marine environments. In 2007, inter-vessel calibrations were conducted to compare catch-per-unit-effort (CPUE) for juvenile salmon between the *John N. Cobb* and the RV *Medeia*.

Calibrations were also done between the *Medeia* and a charter vessel (FV *Steller*) in 2008 and between the *Medeia* and another charter vessel (FV *Chellissa*) in 2009.

From 2010 to 2013, the northern region of Southeast Alaska was sampled monthly from late May to late August with an ABL vessel (May only) and the charter vessel FV *Northwest Explorer*. These surveys included sampling at the Icy Point stations in June, July, and August to complement the Gulf of Alaska Integrated Ecosystem Research Program (GOA-IERP).

Support and rationale for SECM research

In addition to internal NOAA support, the SECM project has been partially supported over the years with funding through the Global Ocean Ecosystem Dynamics (GLOBEC) program, a Pacific Salmon Commission endowment (the Northern Fund, NF), and NOAA's Pacific Coastal Salmon Recovery Fund (Alaska Sustainable Salmon Fund, AKSSF). The SECM research project addresses several goals and objectives identified by the NOAA Science Plan of the AFSC, the NPAFC, and the North Pacific Research Board (NPRB) (GOA-IERP).

The mission of NOAA's current next generation [Strategic Plan](#) involves three objectives: to understand and predict changes in climate, weather, oceans, and coasts; to share that knowledge and information with others; and to conserve and manage coastal and marine ecosystems and resources. NOAA's current vision is to have healthy ecosystems, communities, and economies that are resilient in the face of this change that is expected to impact fisheries production. SECM research emphasizes long-term monitoring of coastal marine habitats used by

juvenile salmon and associated epipelagic fishes, to understand how environmental variation affects the sustainability of these marine resources in an ecological context. SECM research provides valuable time series data specific to salmon and other marine species to monitor long-term ecosystem changes and also provides salmon forecasting data to ensure that regional fisheries are sustained through an ecosystem approach to management. One of the mandates of the [NOAA's Fisheries Priorities and Annual Guidance report for 2014](#) clearly states: "Ensure the productivity and sustainability of fisheries and fishing communities through science-based decision-making and compliance with regulations" and the SECM research project has provided pre-season salmon forecasting information to regional stakeholders of Southeast Alaska fishing communities for the past ten years, thus helping to sustain regional fisheries with science-based forecasts.

The [AFSC Science Plan](#) addresses three main research themes: (1) Monitor and assess fish, crab, and marine mammal populations, fisheries, marine ecosystems, and associated communities that rely on these resources; (2) Understand and forecast effects of climate change on marine resources; and (3) Describe and evaluate the role of habitats in supporting healthy marine ecosystems and populations of fish, crab, and marine mammals. The long-term time series of SECM observations is ideally suited to provide important datasets addressing each of these themes.

The NPAFC 2011–2015 [Science Plan](#) identifies the study of juvenile anadromous salmon stocks in ocean ecosystems as an important component. This Science Plan has a primary goal to explain and forecast the annual variation in Pacific salmon production, an overarching theme of "Forecast of Pacific salmon production in the ocean ecosystems under changing climate", and five primary research topics: 1) Migration and survival mechanisms of juvenile salmon in the

ocean ecosystems; 2) Climate impacts on Pacific salmon production in the Bering Sea (BASIS) and adjacent waters; 3) Winter survival of Pacific salmon in the North Pacific ocean ecosystems; 4) Biological monitoring of key salmon populations; and 5) Development and applications of stock identification methods and models for management of Pacific salmon. SECM research is aligned with both the goal and themes of the NPAFC Science Plan, and specifically addresses research topics 1 and 4 related to the understanding of migration and survival mechanisms of salmon juveniles and biological monitoring of key salmon populations.

Several important components of the NPRB's [GOA-IERP](#) project are addressed by SECM research. These GOA-IERP components include: 1) time series collections of biophysical processes that may provide insight into periodic large-scale regime shifts; 2) carrying capacity and ecological implications of large numbers of hatchery salmon from North America and Asia manifested on forage species in the GOA food web; and 3) input from biological observations including food web shifts of commercially important fish under varying climatic conditions for developing models to forecast annual run strength.

The [U.S. GLOBEC](#) Northeast Pacific program focused on studying “the effects of past and present climate variability on the population ecology and population dynamics of marine biota and living marine resources, and to use this information as a proxy for how the ecosystems of the eastern North Pacific may respond to future global climate change,” and also “...the biophysical mechanisms through which zooplankton and salmon populations respond to physical forcing and biological interactions in the coastal regions....” SECM research addresses the region-scale component of the GLOBEC program by: 1) collecting biological data on juvenile Pacific salmon and ecologically-related fish species from surface rope trawl samples; 2) monitoring physical and biological oceanographic indices at sampling stations in marine

habitats; and 3) conducting process studies focusing on bioenergetics, prey fields, and trophic relationships of juvenile salmon and associated fishes.

Table 1.—Localities and coordinates of stations scheduled for monthly sampling by the Southeast Alaska Coastal Monitoring (SECM) project in marine waters of the northern region of Southeast Alaska in May, June, July, and August of 2014. No sampling is scheduled at the Icy Point stations in May.

Locality	Station	Latitude	Longitude	Offshore distance (km)	Bottom depth (m)
Auke Bay Monitor	ABM	58°22.00'N	134°40.00'W	1.5	60
Upper Chatham Strait	UCA	58°04.57'N	135°00.08'W	3.2	400
Upper Chatham Strait	UCB	58°06.22'N	135°00.91'W	6.4	100
Upper Chatham Strait	UCC	58°07.95'N	135°04.00'W	6.4	100
Upper Chatham Strait	UCD	58°09.64'N	135°02.52'W	3.2	200
Icy Strait	ISA	58°13.25'N	135°31.76'W	3.2	128
Icy Strait	ISB	58°14.22'N	135°29.26'W	6.4	200
Icy Strait	ISC	58°15.28'N	135°26.65'W	6.4	200
Icy Strait	ISD	58°16.38'N	135°23.98'W	3.2	234
Icy Point	IPA	58°20.12'N	137°07.16'W	6.9	160
Icy Point	IPB	58°12.71'N	137°16.96'W	23.4	130
Icy Point	IPC	58°05.28'N	137°26.75'W	40.2	150
Icy Point	IPD	57°53.50'N	137°42.60'W	65.0	1300

Table 2.—Monthly Southeast Alaska Coastal Monitoring (SECM) project research surveys scheduled in marine waters of the northern region of Southeast Alaska in May, June, July, and August of 2014.

Vessel, survey #	On or about (days)	Research focus	Sampling conducted
<i>RV Sashin</i> LV-14-01	20-22 May (3 days)	Oceanography	CTD, chlorophyll and nutrients, zooplankton 9 core stations in Icy/Chatham Straits and Auke Bay
<i>TBD Chartered fishing vessel</i> CV-14-02	26 June–03 July (8 days)	Oceanography Fish survey (trawl)	CTD, chlorophyll and nutrients, zooplankton, fish 9 core stations and 4 stations in Icy Point
<i>TBD Chartered fishing vessel</i> CV-14-03	26 July–02 August (8 days)	Oceanography Fish survey (trawl)	CTD, chlorophyll and nutrients, zooplankton, fish 9 core stations and 4 stations in Icy Point
<i>TBD Chartered fishing vessel</i> CV-14-04	28 August–04 Sept. (8 days)	Oceanography Fish survey (trawl)	CTD, chlorophyll and nutrients, zooplankton, fish 9 core stations and 4 stations in Icy Point




Figure 1.—Stations scheduled for monthly sampling by the Southeast Alaska Coastal Monitoring (SECM) project in marine waters of the northern region of Southeast Alaska in May (no Icy Point stations), June, July, and August of 2014.

Selected publications

- Briscoe, R. J., M. D. Adkison, A.C. Wertheimer, and S. G. Taylor. 2005. Biophysical factors associated with the marine survival of Auke Creek, Alaska, coho salmon. *Trans. Am. Fish. Soc.* 134: 817-828.
- Brodeur, R., E. A. Daly, M. V. Sturdevant, T. W. Miller, J. H. Moss, M. E. Thiess, M. Trudel, L. A. Weitkamp, J. Armstrong, and E. Norton. 2007. Regional comparisons of juvenile salmon feeding ecology in coastal marine waters off the west coast of North America. *Amer. Fish. Soc. Symp.* 57:183-203.
- Fergusson, E. A., M. V. Sturdevant, and J. A. Orsi. 2010. Effects of starvation on energy density of juvenile chum salmon captured in the marine waters of southeastern Alaska. *Fish. Bull.* 108(2): 218–225.
- Fergusson, E. A., M. V. Sturdevant, and J. A. Orsi. 2013. Trophic relationships among juvenile salmon during a 16-year time series of climate variability in Southeast Alaska NPAFC Tech. Rep. 9:112-117. (Available at www.npafc.org).
- Fisher, J. M. T., A. Ammann, J. A. Orsi, J. Piccolo, C. Bucher, E. Casillas, J. A. Harding, B. MacFarlane, R. D. Brodeur, J. F. T. Morris, and D. W. Welch. 2007. Comparisons of the coastal distributions and abundances of juvenile Pacific Salmon from Central California to the Northern Gulf of Alaska. *Amer. Fish. Soc. Symp.* 57:31–80.
- Heard, W. R., J. A. Orsi, A. C. Wertheimer, M. V. Sturdevant, J. M. Murphy, D. G. Mortensen, B. L. Wing, and A. G. Celewycz. 2001. A synthesis of research on early marine ecology of juvenile Pacific Salmon in Southeast Alaska. Page 52 in NPAFC Technical Report 2: Workshop on Factors Affecting Production of Juvenile Salmon. North Pacific Anadromous Fish Commission Tokyo, Japan.
- LaCroix J. J., A. C. Wertheimer, J. A. Orsi, M. V. Sturdevant, E. A. Fergusson, and N. A. Bond. 2009. A top-down survival mechanism during early marine residency explains coho salmon year-class strength in southeast Alaska. *Deep-Sea Research II* 56:2560-2569.
- Mundy, P.R., D.M. Allen, J.L. Boldt, N.A. Bond, S. Dressel, E.V. Farley, Jr., D.H. Hanselman, J. Heifetz, R.R. Hopcroft, M.A. Janout, C. Ladd, R.C. Lam, P.A. Livingston, C.R. Lunsford, J.T. Mathis, F.J. Mueter, C.N. Rooper, N. Sarkar, S.A. K. Shotwell, M. V. Sturdevant, A. C. Thomas, T. J. Weingartner, and D. Woodby. 2010. Status and trends of the Alaska Current Region, 2003-2008, pp. 142-195 In: S.M. McKinnell and M.J. Dagg [Eds.]. *Marine Ecosystems of the North Pacific Ocean, 2003-2008*. PICES Special Publication 4, 393 pp.
- Murphy, J. M., A. L. J. Brase, and J. A. Orsi. 1999. An ocean survey of juvenile salmon in the northern region of southeastern Alaska, May–October 1997. NOAA Tech. Memo NMFS-AFSC-105. Auke Bay Lab., Alaska Fish. Sci. Cen., Nat. Mar. Fish. Serv., NOAA, U.S. Dept. Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA, 40 pp.

- Murphy, J. M., and J. A. Orsi. 1999. Physical oceanographic observations collected aboard the NOAA Ship *John N. Cobb* in the northern region of southeastern Alaska, 1997 and 1998. NOAA Proc. Rep. AFSC 99-02. Auke Bay Lab., Alaska Fish. Sci. Cen., Nat. Mar. Fish. Serv., NOAA, U.S. Dept. Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA, 239 pp.
- Nishimura, G., and M. Sturdevant. 2011. Portable winch technology for use on smaller vessels. *Sea Technology* 52 (July), 3 pp.
- NOAA Strategic Plan 2006-2011. New Priorities for the 21st Century (accessed 11 Jan 2011 at: http://www.nssl.noaa.gov/aboutnssl/Strategic_Plan_2006_FINAL_04282005.pdf).
- NOAA's Next Generation Strategic Plan. 2010. (accessed 20 Jan 2011 at: http://www.ppi.noaa.gov/NGSP3/NGSP_ExecSumm.pdf).
- NPAFC Science Plan 2011-2015 (accessed 11 Jan 2011 at: http://www.npafc.org/new/science_plan.html).
- Orsi, J. A., D. G. Mortensen, and J. M. Murphy. 1999. Early marine ecology of pink and chum salmon in southeastern Alaska. pp 64–72 *In*: Proceedings of the 19th Northeast Pacific Pink and Chum Salmon Workshop, March 3–5 1999, Juneau, AK. Auke Bay Lab., Alaska Fish. Sci. Cen., Nat. Mar. Fish. Serv., NOAA, U.S. Dept. Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA.
- Orsi, J. A., M. V. Sturdevant, J. M. Murphy, D. G. Mortensen, and B. L. Wing. 2000. Seasonal habitat use and early marine ecology of juvenile Pacific salmon in southeastern Alaska. *N. Pac. Anadr. Fish Comm. Bull. No. 2*:111–122.
- Orsi, J. A., M. V. Sturdevant, J. M. Murphy, D. G. Mortensen, B. L. Wing, A. C. Wertheimer, and W. R. Heard. 2001. Southeast Alaska coastal monitoring for habitat use and early marine ecology of juvenile Pacific salmon. Pp. 38 *In*: R. Beamish, Y. Ishida, V. Karpenko, P. Livingston, and K. Myers (eds.). NPAFC Workshop on Factors Affecting Production of Juvenile Salmon: Comparative Studies on Juvenile Salmon Ecology Between the East and West North Pacific Ocean, October 29, 2000, Tokyo, Japan. North Pacific Anadromous Fish Commission Technical Report 2. Vancouver, Canada.
- Orsi, J. A., A. C. Wertheimer, and S. Nutting. 2004. Comparisons of the size of juvenile pink and chum salmon caught during surface rope and pair trawling in the marine waters of southeastern Alaska, June 2004. (NPAFC Doc. 799) Auke Bay Lab., Alaska Fish. Sci. Cen., Nat. Mar. Fish. Serv., NOAA, 11305 Glacier Highway, Juneau, AK 99801-8626, USA, 10 pp.
- Orsi, J. A., A. C. Wertheimer, M. V. Sturdevant, E. A. Fergusson, D. G. Mortensen, and B. L. Wing. 2004. Juvenile chum salmon consumption of zooplankton in marine waters of southeastern Alaska: A bioenergetics approach to implications of hatchery stock interactions. *Rev. Fish Biol. Fish.* 14:335-359.

- Orsi, J. A., J. A. Harding, S. S. Pool, R. D. Brodeur, L. J. Haldorson, J. M. Murphy, J. H. Moss, E. V. Farley Jr., R. M. Sweeting, J. F. T. Morris, M. Trudel, R. J. Beamish, R. L. Emmett, and E. A. Fergusson. 2007. Epipelagic fish assemblages associated with juvenile Pacific salmon in neritic waters of the California Current and the Alaska Current. *Amer. Fish. Soc. Symp.* 57:105-155.
- Orsi, J. A., A. C. Wertheimer, E. A. Fergusson, and M. V. Sturdevant. 2008. Interactions of hatchery chum salmon with juvenile chum and pink salmon in the marine waters of southeastern Alaska. Pp. 20-24 *In: Proceedings of the 23rd Northeast Pacific Pink and Chum Salmon Workshop*. K. Neely, O. Johnson, J. Hard, L. Weitkamp, and K. Adicks (Rapporteurs). February 19-21, 2008, Bellingham, Washington, 95 pp.
- Orsi, J. A., E. A. Fergusson, M. V. Sturdevant, B. L. Wing, A. C. Wertheimer, and W. R. Heard. 2009. Annual Survey of Juvenile Salmon, Ecologically-Related Species, and Environmental Factors in the Marine Waters of Southeastern Alaska, May–August 2008. NPAFC Doc. 1181. 72 pp. (Available at <http://www.npafc.org>).
- Orsi, J. A., M. V. Sturdevant, E. A. Fergusson, B. L. Wing, and W. R. Heard. 2010. Southeast Alaska Coastal Monitoring (SECM) Survey Plan for 2010. NPAFC Doc. 1224. 14 pp. (Available at www.npafc.org).
- Orsi, J., A. Wertheimer, M. Sturdevant, E. Fergusson, and B. Wing. 2009. Insights From a 12-year biophysical time series of juvenile Pacific salmon in Southeast Alaska: the Southeast Alaska Coastal Monitoring Project (SECM). *AFSC, Quarterly July-Aug-Sept. Feature article* (Available at <http://www.afsc.noaa.gov/quarterly/jas2009/jas09.htm>).
- Orsi, J. A., E. A. Fergusson, M. V. Sturdevant, W. R. Heard, and E. V. Farley, Jr. 2011. Annual Survey of Juvenile Salmon, Ecologically-Related Species, and Environmental Factors in the Marine Waters of Southeastern Alaska, May–August 2010*. NPAFC Doc. 1342, 87 pp. (Available at <http://www.npafc.org>). *(The SECM annual survey reports for the years 1997-2009 are also listed in the NPAFC web link as Docs: 277, 346, 497, 498, 536, 630, 702, 798, 871, 955, 1057, 1110, 1181, and 1280)
- Orsi, J. A. 2012. Oral presentations. Chinook salmon insights from marine ecosystem monitoring in Southeast Alaska: Chinook Salmon [Symposium](#), Anchorage, AK in October 2012 and at a [Seminar](#) at the University of Alaska in Juneau, AK in November 2012.
- Orsi, J. A., E. A. Fergusson, and M. V. Sturdevant. 2012. Recent harvest trends of pink and chum salmon in Southeast Alaska: Can marine ecosystem indicators be used as predictive tools for management? NPAFC Tech. Rep. 8:130-134. (Available at www.npafc.org).
- Orsi, J. A., M. V. Sturdevant, E. A. Fergusson, W. R. Heard, and E. V. Farley, Jr. 2013. Chinook salmon marine migration and production mechanisms in Alaska. NPAFC Tech. Rep. 9:240-243. (Available at <http://www.npafc.org>).

- Orsi, J. A., M. V. Sturdevant, E. A. Fergusson, S. C. Heinl, S. C. Vulstek, J. M. Maselko, and J. E. Joyce. 2013. Connecting the “dots” among coastal ocean metrics and Pacific salmon production in Southeast Alaska, 1997-2012. NPAFC Tech. Rep. 9:260-266. (Available at <http://www.npafc.org>).
- Park, W., M. Sturdevant, J. Orsi, A. Wertheimer, E. Fergusson, W. Heard, and T. Shirley. 2004. Interannual abundance patterns of copepods during an ENSO event in Icy Strait, southeastern Alaska. ICES J. Mar. Sci. 61(4):464-477.
- Park, W., and T.C. Shirley. 2008. Variations of abundance and hatch timing of dungeness crab larvae in southeastern Alaska: implications for climate effect. Animal Cells and Systems 12:287-295.
- Reese C, N. Hillgruber, M. Sturdevant, A. Wertheimer, W. Smoker, and R. Focht. 2009. Spatial and temporal distribution and the potential for estuarine interactions between wild and hatchery chum salmon (*Oncorhynchus keta*) in Taku Inlet, Alaska. Fish. Bull. 107:433–450.
- Robbins, J. B. 2006. Biophysical factors associated with the marine growth and survival of Auke Creek, Alaska, coho salmon (*Oncorhynchus kisutch*). Unpub. M.S. Thesis, University of Alaska, Juneau, Alaska. 93 pp.
- Sturdevant, M. V., E. A. Fergusson, J. A. Orsi, and A. C. Wertheimer. 2002. Diel feeding of juvenile pink, chum, and coho salmon in Icy Strait, southeastern Alaska, May–September 2001 (NPAFC Doc. 631) Auke Bay Lab., Alaska Fish. Sci. Cen., Nat. Mar. Fish. Serv., NOAA, U.S. Dept. Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA, 42 pp.
- Sturdevant, M. V., E. A. Fergusson, J. A. Orsi, and A. C. Wertheimer. 2004. Diel feeding and gastric evacuation of juvenile pink and chum salmon in Icy Strait, Southeastern Alaska, May–September 2001. NPAFC Tech. Rep. No. 5: 107-109. Auke Bay Lab., Alaska Fish. Sci. Cen., Nat. Mar. Fish. Serv., NOAA, U.S. Dept. Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626.
- Sturdevant M. V., E. A. Fergusson, J. A. Orsi, and A. C. Wertheimer. 2008. Seasonal patterns in diel feeding, gastric evacuation, and energy density of juvenile chum salmon in Icy Strait, Southeast Alaska, 2001. *In*: Proceedings of the 23rd Northeast Pacific Pink and Chum Salmon Workshop. K. Neely, O. Johnson, J. Hard, L. Weitkamp, and K. Adicks (Rapporteurs). February 19-21, 2008, Bellingham, Washington, 95 pp.
- Sturdevant, M. V., M. F. Sigler, and J. A. Orsi. 2009. Sablefish predation on juvenile salmon in the coastal marine waters of Southeast Alaska in 1999. Trans. Am. Fish. Soc. 138:675–691.
- Sturdevant, M., G. Nishimura, and J. Orsi. 2011. Sidewinder: description of a new block winch for deploying instruments at sea. Mar. Coastal Fish. 3(1): 317-323.

- Sturdevant, M., E. Fergusson, N. Hillgruber, C. Reese, J. Orsi, R. Focht, A. Wertheimer, and B. Smoker. 2012. Lack of trophic competition among wild and hatchery juvenile chum salmon during early marine residence in Taku Inlet, Southeast Alaska. *Environmental Biology of Fishes* 94:101-116. Available at <http://dx.doi.org/10.1007/s10641-011-9899-7>.
- Sturdevant, M. V., J. A. Orsi, and E. A. Fergusson. 2012. Diets and trophic linkages of epipelagic fish predators in coastal Southeast Alaska during a period of warm and cold climate years, 1997–2011. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 4(1):526-545, Available at: <http://www.tandfonline.com/doi/abs/10.1080/19425120.2012.694838>
- Sturdevant, M. V., R. E. Brenner, E. A. Fergusson, J. A. Orsi, and W. R. Heard. 2013. Does predation by returning adult pink salmon regulate pink salmon or herring abundance? NPAFC Tech. Rep. 9: 153-164. (Available at <http://www.npafc.org>).
- Trudel, M., J. Fisher, J. A. Orsi, J. F. T. Morris, M. E. Thiess, R. M. Sweeting, S. Hinton, E. A. Fergusson, and D. W. Welch. 2009. Distribution and Migration of Juvenile Chinook salmon derived from coded wire tag recoveries along the continental shelf of western North America. *Trans. Amer. Fish. Soc.* 138:1369–1391.
- Turner, E., and D. B. Haidvogel. 2009. [Taking ocean research results to applications: examples and lessons from US GLOBEC](#). *Oceanography* 22(4):232-241.
- Weitkamp, L. A. and M. V. Sturdevant. 2008. Food habits and marine survival of juvenile Chinook and coho salmon from marine waters of Southeast Alaska. *Fish. Oceanog.* 17(5):380–395.
- Weitkamp, L. A., J. A. Orsi, K. W. Myers, and R. C. Francis. 2011. Contrasting early marine ecology of Chinook and coho salmon in Southeast Alaska: insight in to factors affecting marine survival. *Mar. Coastal Fish.* 3(1):233-249.
- Wertheimer, A. C., E. A. Fergusson, R. L. Focht, W. R. Heard, J. A. Orsi, M. V. Sturdevant, and B. L. Wing. 2003. Sea lice infestation of juvenile salmon in the marine waters of the northern region of southeastern Alaska, May–August, 2003. (NPAFC Doc. 706) Auke Bay Lab., Alaska Fish. Sci. Cen., Nat. Mar. Fish. Serv., NOAA, U.S. Dept. Commerce, 11305 Glacier Highway, Juneau, AK 99801-8626, USA, 13 pp.
- Wertheimer, A. C., J. A. Orsi, M. V. Sturdevant, and E. A. Fergusson. 2008. Forecasting pink salmon abundance in Southeast Alaska from juvenile salmon abundance and associated environmental parameters. Final Report, Pacific Salmon Commission Northern Fund. 41 pp.
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2009. Calibration of Juvenile Salmon Catches using Paired Comparisons between Two Research Vessels Fishing Nordic 264 Surface Trawls in Southeastern Alaska, July 2008. NPAFC Doc. 1180. 18 pp. (Available at <http://www.npafc.org>).

- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2009. Forecasting Pink Salmon Harvest in Southeast Alaska from Juvenile Salmon Abundance and Associated Environmental Parameters: 2008 Returns and 2009 Forecast. NPAFC Doc. 1201. 20 pp. (Available at <http://www.npafc.org>).
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2010. Calibration of Juvenile Salmon Catches using Paired Comparisons between Two Research Vessels Fishing Nordic 264 Surface Trawls in Southeast Alaska, July 2009. NPAFC Doc. 1277, 20 pp. (Available at <http://www.npafc.org>).
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2010. Forecasting Pink Salmon Harvest in Southeast Alaska from Juvenile Salmon Abundance and Associated Environmental Parameters: 2009 Harvest and 2010 Forecast. NPAFC Doc. 1278, 20 pp. (Available at <http://www.npafc.org>).
- Wertheimer, A. C., J. A. Orsi, E. A. Fergusson, and M. V. Sturdevant. 2011. Forecasting Pink Salmon Harvest in Southeast Alaska from Juvenile Salmon Abundance and Associated Environmental Parameters: 2010 Returns and 2011 Forecast. NPAFC Doc. 1343, 20 pp. (Available at <http://www.npafc.org>).