

Canadian Salmon Catch and Enhanced Salmon Production in 2017 and 2018

by

Mary E. Thiess¹, Joan Bateman², Shelee Hamilton¹, and Mark Beere³

¹Fisheries and Oceans Canada
3190 Hammond Bay Road
Nanaimo, B.C., Canada V9T 6N7

²Fisheries and Oceans Canada
200-401 Burrard Street
Vancouver, B.C., Canada V6C 3S4

³B.C. Ministry of Forests, Lands and Natural Resource Operations
Fisheries Branch—Skeena, Bag 5000
Smithers, B.C., Canada V0J 2N0

Submitted to the

NORTH PACIFIC ANADROMOUS FISH COMMISSION

by

Canada

April 2019

THIS PAPER MAY BE CITED IN THE FOLLOWING MANNER:

Thiess, M.E., J. Bateman, S. Hamilton, and M. Beere. 2019. Canadian salmon catch and enhanced salmon production in 2017 and 2018. NPAFC Doc. 1837. 9 pp. Fisheries and Oceans Canada, and B.C. Ministry of Forests, Lands and Natural Resource Operations (Available at <https://npafc.org>).

ABSTRACT

This document reports final catch estimates for 2017 and preliminary catch estimates for 2018 for the six major salmon species in British Columbia (B.C.) and Yukon fisheries. Catch is reported for commercial fisheries (numbers and total weight) in tidal waters, and recreational (numbers only) and aboriginal fisheries (numbers only) in tidal and non-tidal waters. Catches include non-Canadian origin fish caught in B.C. and exclude Canadian origin fish caught in fisheries outside B.C. This document also summarizes release information for salmon including steelhead trout from Fisheries and Oceans Canada (DFO) and Freshwater Fisheries Society of B.C. enhancement facilities in B.C. in 2017 and 2018.

Keywords: Salmon, steelhead, catch, enhancement, British Columbia

INTRODUCTION

The six species of salmon native to B.C. contribute to commercial, recreational, and aboriginal fisheries. Commercial net fisheries tend to focus on sockeye (*Oncorhynchus nerka*), chum (*O. keta*), and pink (*O. gorbuscha*) salmon, while recreational harvesters catch mostly Chinook (*O. tshawytscha*), coho (*O. kisutch*), and steelhead (*O. mykiss*). All species of salmon are valued by aboriginal fisheries, which typically occur in terminal areas within freshwater, but harvest also occurs in marine waters. Steelhead trout constitute a relatively minor component of the catch.

This document reports final catch estimates for 2017 and preliminary 2018 catch estimates of Pacific salmon in B.C. and Yukon fisheries. Estimates are reported for retained commercial catch (numbers and total weight) in tidal waters (except the Yukon fisheries) and recreational (numbers only) and aboriginal catch (numbers only) in tidal and non-tidal waters by species. Catches include non-Canadian origin fish caught in B.C. and exclude Canadian origin fish caught outside B.C.

The objectives of the DFO Salmonid Enhancement Program (SEP) are to rebuild stocks and increase catch through the expanded use of enhancement technology. The program produces Chinook, coho, chum, pink, and sockeye salmon, as well as small numbers of steelhead and cutthroat trout (*O. clarki*). Steelhead and cutthroat enhancement takes place at SEP facilities under a cooperative arrangement with the Government of B.C. Projects include hatcheries, fishways, spawning and rearing channels, and small classroom incubators; releases range in size from nearly 100 million juveniles from spawning channels, to less than one hundred from school classroom projects, annually.

CATCH ESTIMATES

A brief description of the data and sources of information for the four major fishery types follows:

Commercial Catch Estimates

The Fisheries Operating System (FOS) is the official salmon commercial fishery catch database for DFO in the Pacific Region. FOS maintains various fisher-produced data (e.g. sale slips,

logbooks, at-sea interviews) and fisher-independent data (e.g. on-board observers and dockside monitors). The final commercial catch estimates reported here for 2017 and 2018 are manager-determined estimates based on resolution of all data sources.

Commercial catch estimates for 2017 and 2018 are presented as numbers and weights (Table 1) of fish retained by commercial salmon vessels (salmon gill net, salmon seine net and salmon troll). Weights are based on sale slip data and are given as round (i.e., whole fish) equivalents. Sale slips record gear type, area of catch, date of landing, and landed weight and value by species. Since the value of the landing and resulting payments are based primarily on the landed weight, weight estimates are considered reasonably accurate (Sandher et al. 2012).

Piece estimates recorded on sale slips are usually estimated by applying an estimated average weight to the landed weight, particularly for high volume net landings (i.e., net landings of sockeye, pink and/or chum salmon). The average weight estimates are obtained by weighing counted samples of landed catch during offloading. Sampling for average weights is conducted to allow relatively accurate and precise, species-, area-, time- and gear-specific estimates of average weights to be generated. Such estimates have been widely available for Chinook and coho salmon since the early 1980s and for sockeye, chum and pink salmon since 1994.

Recreational Catch Estimates (non-Steelhead)

Recreational salmon catch estimates (non-Steelhead) for 2017 and 2018 are reported in pieces (Table 2). Creel surveys are the primary source of recreational catch data, resulting in estimates by DFO Pacific Fishery Management Area (PFMA) and month. Creel surveys generally cover the times and areas with relatively large effort and catch, but typically not times and areas where fishing is open with relatively low effort and/or catch rates. Catch from sport fishing lodges are based on logbook programs where they are in use. Catch from lodges not reporting via logbooks is incorporated in creel survey-based estimates. Further, the numbers provided here are the unexpanded estimates from the creel and logbook programs; they do not include estimates for catch from times and areas not surveyed. Consequently, recreational catches reported here underestimate total catch.

Recreational Steelhead Catches—A Historical Overview

Although there are no targeted fisheries on steelhead in marine waters, valuable freshwater recreational fisheries occur and are a provincial responsibility delegated from the Government of Canada. The B.C. management objectives for these fisheries are:

1. Maintain a diversity of sustainable recreational angling opportunities for steelhead in British Columbia.
2. Maintain, protect and restore the productive capacity of the freshwater environment to produce steelhead (B.C. Ministry of Forests, Lands and Natural Resource Operations, July 2014).

There has been no retention of wild steelhead in B.C. sport fisheries since April 1, 2007; recreational fisheries that currently operate on non-enhanced rivers are strictly catch and release. Recreational fisheries also occur in approximately 13 hatchery-augmented rivers, where an angler is allowed to harvest one fish per day (with a maximum 10 per season). Total catches are estimated via the Steelhead Harvest Analysis – a questionnaire mailed to 50% of all B.C. and 100% of non-B.C. residents who purchased a steelhead licence. Estimates are currently available

up to the end of the 2017/2018 fiscal year (M. Beere, B.C. Ministry of Forests, Lands and Natural Resource Operations, Smithers; pers. comm., March 12, 2019; Table 3). Conservation concerns about two populations of Interior Fraser River steelhead led to emergency listings as “endangered” in January 2018. It is expected that actions taken as a result of this listing will impact future catches of steelhead in B.C. waters.

Aboriginal Catch Estimates

Aboriginal (subsistence) catch represents retained catch and is reported in pieces (Table 4). Aboriginal catch statistics can usually be resolved to PFMA and year, but areas may be grouped in some cases. A variety of approaches have been used to generate estimates. While aboriginal catch estimation programs cover many of the largest aboriginal fisheries, coverage is incomplete both temporally and spatially. Similar to recreational catch, aboriginal catch estimates presented here do not include catch from times and areas not covered by catch estimation programs, and therefore underestimate total catch.

ENHANCED PRODUCTION

Annual egg and juvenile stock production targets for hatcheries are set pre-season, in consultation with project managers, stock assessment biologists and harvest management biologists. Production objectives include conservation or rebuilding, harvest and/or assessment. Other considerations include potential species and stock interactions, effects on natural stocks, harvest concerns, habitat capacity and project capacity. The production plan is finalized after review by industry, the public and other interested groups in the Salmon Integrated Fisheries Management Planning process.

Depending on the species and enhancement approach, juvenile fish are released at various life stages. Chum and pink salmon are released either immediately after emergence as unfed fry or after one month of feeding (fed fry). Coho are released as fry, either at emergence or after three to five months of rearing, or as smolts after one year of rearing. In B.C. sockeye enhancement is typically conducted using spawning and rearing channels rather than hatcheries, and juveniles emigrate volitionally soon after emergence. Sockeye are also enhanced through lake fertilization programs, and most migrate to sea after one year of lake rearing. Coastal ocean-type chinook stocks are released after three to four months of rearing. Interior stream-type stocks are frequently reared for one year, and generally constitute a very small component of the total numbers of chinook released. Releases from hatcheries are usually estimated by subtracting known egg and fry mortalities from egg numbers. Releases from rearing channels are estimated by sampling out-migrants.

Release information for salmon from DFO enhancement facilities and the Fraser Valley Trout Hatchery (Freshwater Fisheries Society) in B.C. in 2017 (final) and 2018 (preliminary) is summarized in Table 5. Approximately 360.6 and 261.7 million salmon were released, in 2017 and 2018, respectively. In both years, sockeye and chum releases accounted for the majority of enhanced production (84% and 77% of total releases in 2017 and 2018, respectively). Since 2016, all thirteen hatcheries augmenting steelhead production have conducted smolt release programs, releasing approximately 354,000 and 285,000 steelhead smolts in 2017 and 2018, respectively (Table 5).

It is not possible to conduct assessments of each enhancement project and/or release strategy individually. Consequently, certain stocks are used as indicators, their production is tagged or marked annually and rigorous fishery and escapement sampling and estimation programs are conducted. Survival and exploitation estimates are used for time series analyses of both wild and enhanced populations. Enhanced contributions and survivals of chinook, coho, and chum salmon are normally estimated by applying a tag and /or external mark to a portion of the fish released and subsequently recovering these marked fish through sampling programs in the fisheries, on the spawning grounds and at enhancement sites. Visual marks are chiefly the removal of the adipose fin, with some chinook and coho stocks also receiving an implanted coded wire tag to enable identification of stock and release year. Beginning in 1996, most enhanced coho from southern B.C. have been marked with a fin clip to enable the prosecution of mark-selective fisheries. Some species and stocks are given a thermal otolith mark to estimate enhanced contributions to terminal areas. Thermal marking is coordinated with the DFO Salmon Assessment Section and is reported separately.

ACKNOWLEDGEMENTS

Many thanks to the following individuals for their updates and analysis of the data included in this report: Karen Burnett, Denise Clark, Kevin Conley, Kris Hein, Jeff Radford, Jamie Scroggie, Joe Tadey, and Bill Waugh.

REFERENCES

- Ministry of Forests, Lands and Natural Resource Operations (FLNRO). 2014. Provincial Framework for Steelhead Management in British Columbia. Fish, Wildlife and Habitat Management Branch, Victoria, B.C. DRAFT Report, July 2014.
http://www.bcfdf.com/bcfdf_news/Framework.pdf
- Sandher, J., C. Lynch, D. Willis, R. Cook and J. R. Irvine. 2012. Canadian enhanced salmonid production during 1977–2011 (1976–2010 brood years). NPAFC Doc. 1420. 10 pp. Fisheries and Oceans Canada. (Available at <https://npafc.org>).

Table 1. Canadian commercial catch estimates of Pacific salmon (excluding steelhead) for 2017 and 2018, presented in pieces (000's) and weights (MT). Catches reported include First Nations commercial fisheries.

Reporting Area	Year	Commercial Catch Number (000's)						Commercial Catch Round Weight (MT)					
		Pink	Chum	Sockeye	Coho	Chinook	Total	Pink	Chum	Sockeye	Coho	Chinook	Total
Fraser River	2017	4.13	197.59	29.85	0.73	1.74	234.03	9.32	967.96	74.32	2.20	10.91	1,064.71
	2018	0.04	9.17	1,730.93	1.45	2.29	1,743.88	0.08	45.54	4,647.37	4.33	11.60	4,708.92
South Coast	2017	37.24	760.23	11.00	0.46	0.06	808.99	1.9	82.80	3,664.34	27.42	1.51	3,776.47
	2018	41.51	211.33	655.93	0.08	0.06	908.91	81.1	76.00	942.97	1,733.72	0.24	2,753.20
West Coast Vancouver Island	2017	0.03	56.65	56.85	2.71	99.00	215.23	0.05	270.29	114.05	9.61	571.45	965.45
	2018	0.02	27.56	71.26	6.00	73.81	178.66	0.05	126.93	160.78	23.67	311.91	623.33
Central Coast	2017	748.33	333.26	0.77	8.35	3.15	1,093.86	1,662.14	1,860.06	1.96	24.33	21.41	3,569.90
	2018	48.48	452.44	3.56	0.00	4.60	509.09	107.90	2,423.27	9.30		20.83	2,561.31
North Coast	2017	835.92	77.32	50.38	84.75	1.93	1,050.29	1,854.52	396.36	125.81	262.48	10.96	2,650.13
	2018	122.15	46.96	120.60	35.55	0.51	325.76	267.67	235.16	278.02	102.99	3.00	886.84
Haida Gwaii	2017	33.06	0.34		273.19	97.59	404.18	55.58	1.49	0.00	816.51	655.82	1,529.39
	2018	27.97	3.18		142.19	70.28	243.62	62.58	14.52	0.01	411.62	472.18	960.91
Yukon/ Transboundary	2017		2.40	63.10	13.20	1.30	80.00	0.00	8.49	154.56	54.49	7.84	225.38
	2018	0.10	0.53	35.27	13.19	0.00	49.09	0.13	1.89	69.83	43.56	0.00	115.41
Whole Country	2017	1,658.69	1,427.78	211.96	383.38	204.78	3,886.59	3,664.41	7,169.00	498.12	1,171.13	1,278.78	13,781.44
	2018	240.27	751.17	2,617.55	198.46	151.56	3,959.00	514.41	3,790.28	6,899.03	586.40	819.80	12,609.92

Table 2. Canadian recreational catch estimates of Pacific salmon (excluding steelhead) for 2017 and 2018, presented in pieces (000's).

Reporting Area	Year	Recreational Catch Numbers (000's)						Total
		Pink	Chum	Sockeye	Coho	Chinook		
South Coast	2017	29.34	5.21	0.24	29.32	111.90	176.02	
	2018	4.73	5.85	159.10	44.35	124.69	338.72	
West Coast Vancouver Island	2017	2.43	0.03	11.45	27.04	95.21	136.16	
	2018	0.07	0.01	5.71	45.86	81.84	133.49	
North Coast	2017	3.96	0.27	0.29	58.50	18.03	81.06	
	2018	1.39	0.18	0.03	10.44	5.82	17.86	
Haida Gwaii	2017	1.15	0.90	0.17	35.10	45.60	82.92	
	2018	1.95	0.95	0.17	34.20	36.70	73.97	
Yukon/Transboundary	2017	0.0	0.0	0.1	0.0	0.2	0.3	
	2018	NR	NR	NR	NR	NR	NR	
Whole Country	2017	36.88	6.41	12.25	149.97	270.95	476.45	
	2018	8.14	6.98	165.01	134.85	249.06	564.04	

Table 3. Canadian recreational catch estimates of steelhead since 2005/2006, presented in pieces (000's). Total catches are estimated via the Steelhead Harvest Analysis (FLNRO 2014). B.C. Regions are defined as follows: 1=Vancouver Island; 2=Lower Mainland; 3=Thompson; 4=Kootenay; 5=Cariboo; 6=Skeena. Data provided by M. Beere, B.C. Ministry of Forests, Lands and Natural Resource Operations, Smithers.

Year	B.C. Region					Unknown	Annual Total
	1	2	3	5	6		
2005/2006	16.842	22.624	2.730	2.635	32.311	0.259	77.401
2006/2007	22.330	11.466	1.813	3.698	25.366	0.372	65.045
2007/2008	15.929	14.031	0.809	2.491	26.951	0.358	60.569
2008/2009	0.000	0.000	0.000	0.000	0.000	0.000	0.000
2009/2010	20.137	9.637	1.204	2.017	38.790	0.579	72.364
2010/2011	18.021	23.400	0.023	4.389	45.265	0.370	91.468
2011/2012	16.592	27.163	1.633	5.230	36.311	0.235	87.164
2012/2013	14.123	25.681	1.922	6.777	55.479	0.333	104.315
2013/2014	8.294	19.299	2.005	3.693	41.895	0.501	75.687
2014/2015	13.003	23.603	1.361	5.341	51.240	0.769	95.317
2015/2016	8.100	14.191	0.244	8.027	43.237	0.525	74.324
2016/2017	6.259	11.149	0.561	5.530	44.081	0.854	68.434
2017/2018	8.042	10.196	0.141	3.600	24.050	0.175	46.204

Table 4. Canadian subsistence catch estimates of Pacific salmon (including steelhead) for 2017 and 2018, presented in numbers (000's).

Reporting Area	Year	Subsistence Numbers (000's)						Total
		Pink	Chum	Sockeye	Coho	Chinook	Steelhead	
South Coast	2017	46.35	82.68	71.65	16.59	30.56		247.83
	2018	3.94	78.15	872.87	1.82	36.35		993.13
West Coast Vancouver Island	2017	0.02	54.91	24.67	14.15	49.96		143.71
	2018		41.654	17.967	10.346	31.791		101.76
North Coast	2017	17.46	2.52	116.30	13.52	11.67	0.89	162.36
	2018	3.82	0.87	129.91	5.59	10.91	3.67	154.77
Haida Gwaii	2017			4.87				4.87
	2018	5.00	4.00	7.17		1.00		17.17
Yukon/Transboundary	2017		3.30	9.40	0.80	1.10		14.60
	2018		2.87	5.43	0.032	3.27		11.60
Whole Country	2017	63.83	143.41	226.89	45.07	93.29	0.89	573.37
	2018	12.76	127.54	1,033.35	17.79	83.32	3.67	1,278.44

Table 5. DFO Salmon Enhancement Program and Fraser Valley Hatchery enhanced Pacific salmon releases (including steelhead) in 2017 and 2018, presented in numbers (000's).

Reporting Area	Release Year	Enhanced Salmon Releases (000's)						Total
		Pink	Chum	Sockeye	Coho	Chinook	Steelhead	
B.C. Interior	2017			14,902	174	1,206		16,283
	2018			3,276	350	1,223		4,849
Fraser River	2017		9,382	1,483	1,786	2,229	193	15,074
	2018	526	13,274	25,729	1,722	2,249	198	43,698
South Coast	2017	11,044	82,156	257	3,771	14,428	68	111,724
	2018	14,195	27,780	530	4,043	15,006	48	61,602
West Coast Vancouver Island	2017		30,968		963	15,899	41	47,871
	2018		11,124		843	15,270	0	27,237
Central Coast	2017		9,257	133	30	2,429		11,849
	2018		8,619	63	73	2,363		11,118
North Coast	2017		1,767	152,634	682	2,224	52	157,358
	2018		883	109,063	842	1,995	39	112,822
Haida Gwaii	2017		131		196	88		415
	2018		155		129	118		402
Whole Country	2017	11,044	133,661	169,409	7,603	38,503	354	360,574
	2018	14,721	61,835	138,661	8,002	38,224	285	261,728