Interannual variations of prey resources during the early ocean life of juvenile chum salmon in four coastal areas around Hokkaido, northern Japan

Tomoki SATO, Kei SASAKI, Masaya TAKAHASHI and Toru NAGASAWA
(Hokkaido National Fisheries Research Institute, Fisheries Research Agency.)

Introduction

The early life of juvenile chum salmon is a critical phase in their life history. It is believed high mortality occurs in this initial phase, which may be an important determinant of subsequent salmon population size. The state of prey environment such as zooplankton abundance and species composition is important for the growth and survival of juvenile chum salmon during their early ocean life. In order to evaluate prey environment during the early ocean life of juvenile chum salmon between March and July, 1998-2010, we conducted a long-term zooplankton monitoring in four coastal areas around Hokkaido, northern Japan. We will report interannual variations of prey environment for juvenile chum salmon during early ocean life in the each areas.

Materials & methods

Zooplankton were collected with vertical tows by a NORPAC net (0.45 m in mouth diameter, 0.33 mm mesh) from 20 m to surface at each site. If the water depth less than 20 m, the net was towed from 3 m above bottom to surface.

Fig. 1. Location of sampling area

Table 1. Sampling station, distance from seacoast (km) and water depth (m) at each sampling areas.

Areas	Stn. No	Distance from seacoast (km)	Depth (m)
ATSUTA Japan Sea coast	Stn. 1	0.5	10
	Stn. 2	1.0	15
	Stn. 3	2.0	20
	Stn. 4	10.0	30
SHIRAOI Western Pacific coast	Stn. 1	0.5	7
	Stn. 2	1.0	10
	Stn. 3	1.5	20
	Stn. 4	2.0	30
KONBUMORI Eastern Pacific coast	Stn. 1	0.4	15
	Stn. 2	1.3	21
	Stn. 3	3.5	40
	Stn. 4	7.8	60
SHARI Okhotsk Sea	Stn. 1	1.0	10
	Stn. 2	2.0	19
	Stn. 3	3.7	38
	Stn. 4	5.4	48

Table 2. List of occurrence species in the for coastal areas during the period from 1998 to 2010.

Hydrozoans	Molluscs	Polychaetes	Cladocerans	Calanoid copepods (<2 mm TL)	Calanoid copepods (> 2mm TL)	Other copepods
Haliscera sp.	Gastropod larvae	Unidentified larvae	Podon leuckartl	Acartia hudsonica	Acartia longiremis	Oithona atlantica
Sarsia sp.	Bivalve larvae		Podon polyphemoides	Calanus glacialis	Acartia omorii	Oithona similis
Rathkea octopunctata			Evadne nordmanni	Calanus pacificus	Acartia steueri	Oithona spp. (nauplius)
Dataria sp.			Cercopus rotundi	Calanus rotundi	Eucalanus bungii	Oithona sp. (copepodite)
Aglantha sp.			Evadne spinifera	Eucalanus bungii	Metridia okhotensis	Microsetella norvegica
Diphyes sp.				Metridia pacifica	Neocalanus cristatus	Microsetella rosea
				Metridia okhotensis	Neocalanus plumchrus	Oncaea mediterranea
				Neocalanus cristatus	Neocalanus flemingeri	Corycaeus affinis
				Neocalanus flemingeri	Neocalanus plumchrus	Oncaea mediterranea
				Neocalanus plumchrus	Pseudocalanus minutus	Corycaeus lubbocki
				Pseudocalanus minutus	Pseudocalanus moultoni	Corycaeus sp.
				Scolecithricella minor	Eurytemora pacifica	
				Tortanus discaudatus	Eurytemora herdmani	
					Centropages abdominalis	
					Centropages tenuiremis	
					Epilabidocera longipedata	
					Metridia pacifica	
					Metridia okhotensis	
					Neocalanus cristatus	
					Neocalanus flemingeri	
					Neocalanus plumchrus	
					Pseudocalanus minutus	
					Pseudocalanus moultoni	
					Scolecithricella minor	
					Tortanus discaudatus	

Cirripeds	Amphipods	Euphausiids
Balanomorpha (nauplius)	Hyperia galba	Euphausia pacifica (furcilia)
Balanomorpha (cypris)	Themisto japonica	Thysanoessa spp. (furcilia)
Pollicipes polymerus (nauplius)	Gammaridae (unidentified)	Euphausiacea (egg)
	Hyperiidea (unidentified)	Euphausiacea (nauplius)
	Caprella sp.	Euphausiacea (metanauplius)
		Euphausiacea (calyptopis)

Decapods	Other crustaceans	Chaetognaths	Echinoderms	Appendicularians	Fishes	Others
Carides (zoea)	Mysidacea	Sagitta elegans	Asteroidea	Oikopleura dioica	Fish eggs (Engraulis japonicus)	Aphelascolex larvae
Thalassinidea (zoea)	Cumacea	Sagitta spp.	Ophiuroidea	Oikopleura albicans/vanhoffeni	Fish eggs (unidentified)	Cyphonautes larvae
Anomura (zoea)	nauplius	Sagittidae (juvenile)	Echinoidea	Oikopleura longicauda	Fish larvae (unidentified)	Pilidium larvae
Brachyura (zoea)		Sagittoidea (eggs)	Ophiuroidea (pluteus)	Oikopleura spp. (juvenile)		
				Fritillaria borealis f.typica		

ATSUTA (Japan Sea coast)



Fig. 2. Change in mean water temperature (April – June) of 3 m depth at stn. 3 (a), abundance (mean±SE) of total zooplankton (b), and composition of top 10 dominant species (c) during the period from 1998 to 2010 in the Atsuta area.



Fig. 3. Change in abundance of zooplankton species mainly found in stomach of juvenile chum salmons the Atsuta area. Error bars are standard error of the means.

SHARI (Okhotsk Sea coast)



Fig. 6. Change in mean water temperature (April – June) of 3 m depth at stn. 2 (a), abundance (mean±SE) of total zooplankton (b), and composition of top 10 dominant species (c) during the period from 1998 to 2010 in the Shari area.



Fig. 7. Change in abundance of zooplankton species mainly found in stomach of juvenile chum salmons the Shari area. Error bars are standard error of the means.

SHIRAOI (Western Pacific coast)



Fig. 4. Change in mean water temperature (April – June) of 3 m depth at stn. 3 (a), abundance (mean±SE) of total zooplankton (b), and composition of top 10 dominant species (c) during the period from 1998 to 2010 in the Shiraoi area.



Fig. 5. Change in abundance of zooplankton species mainly found in stomach of juvenile chum salmons the Shiraoi area. Error bars are standard error of the means.

KONBUMORI (Eastern Pacific coast)



Fig. 8. Change in mean water temperature (April – June) of 3 m depth at stn. 2 (a), abundance (mean±SE) of total zooplankton (b), and composition of top 10 dominant species (c) during the period from 1998 to 2010 in the Konbumori area.



Fig. 9. Change in abundance of zooplankton species mainly found in stomach of juvenile chum salmons the Konbumori area. Error bars are standard error of the means.

Conclusion

Dominant zooplanktons were small copepods such as *Pseudocalanus newmani* in Japan Sea and Okhotsk Sea coast, and *Acartia longiremis* in the Pacific coasts. However, juvenile chum salmon fed larger zooplanktons more than 2 mm in body length. A trend of interannual variation of prey organisms was different in four coastal areas around Hokkaido.

Future works

We need to evaluate how interannual variations of abundance of prey organisms affect the growth and survival of juvenile chum salmon in each coastal waters.