

Russian National Overview of BASIS Research

Vyacheslav P. Shuntov¹, Olga S Temnykh¹, Vladimir V. Sviridov¹, and Vladimir I. Karpenko²

¹Pacific Scientific Research Fisheries Centre (TINRO-Centre),
4, Shevchenko Alley, Vladivostok 690950, Russia

²Kamchatka Fishery and Oceanography Research Institute (KamchatNIRO),
18, Nabereznaya Street, Petropavlovsk-Kamchatsky 683000, Russia



Keywords: Nekton community, salmon, biomass, distribution

BASIS research by the R/V *TINRO* was conducted in the fall of 2002 (86 sampling stations) and summer and fall of 2003 (176 sampling stations) in the western Bering Sea and adjacent Pacific waters. Each sampling station included a trawl tow, as well as, hydrobiological and hydrological stations, which were carried out in accordance with standard methods of TINRO-Centre (Temnykh et al., 2004). The results of Russian research within BASIS program (2002 and 2003) are as follows:

1. In 2002 and 2003 the tendency of hydrological processes in the western Bering Sea followed a warm year type. Intensification of water exchange between the Bering Sea and Pacific Ocean was accompanied with the rise of SST and the increase of its discharge through the Kamchatsky Strait (Khen and Basyuk 2005).
2. During the last two years, a marked decrease in plankton biomass through the reduction of copepods abundance was noted for Aleutian and Commander basins within the Russian EEZ. On the other hand, relative abundance of euphausiids has increased (Volkov et al. 2005).
3. Total nekton biomass in the western Bering Sea was estimated as 1.7–2.8 million tons in fall of 2002–2003 and 1.3 million tons in summer of 2003. In the summer of 2003 the biomass of Pacific salmon exceeded 939 thousand tons. During summer-autumn period early 2000-ies a significant increase of Pacific salmon share in total fish biomass in upper epipelagic layer took place (Fig. 1). The biomass of salmon distributed in shelf zone was 10–100 times lower than their abundance in the deeper-water within the Bering Sea basin (Fig. 2) (Temnykh, 2004). According to the diel study at different water strata the major concentrations of salmon are located in the upper 30 m.
4. Comparative analysis of spatio-temporal variability of Pacific salmon's biological characteristics suggests species-specific and age-specific adaptive strategy, which is aimed at the lessening of density-dependent interactions and maximum utilization of available feeding grounds (Sviridov et al. 2004a).
5. Asian stocks of immature age .1 and .2 sockeye salmon dominate within the southwestern Bering Sea and the adjacent waters of the northwest Pacific Ocean in September–October (Bugaev 2003, 2004a, 2004b). There were differences in the interspecies structure in the catches for different age groups that reflects divergence in distribution range among different age regional groupings of salmon during their marine growing period.
6. The rates of different types of traumatization and infestation occurrence are species- and age-specific and are subject to significant spatio-temporal variability (Sviridov et al. 2004b).

Fig. 1. Interannual dynamics of fish communities composition (% of total biomass) in the upper epipelagic layer in the southwestern Bering Sea (within Russian EEZ) in summer 1991–2003.





Fig. 2. Ratio (%) of Pacific salmon biomass in shelf and deep-water regions of the western Bering Sea.



7. Analyses of long-term datasets on salmon food habits in the western Bering Sea suggests that the composition and amount of Pacific salmon ration in 2000s is not significantly different between the 1980s to mid-1990s period (Temnykh 2004; Temnykh *et al.* 2004; Shuntov and Temnykh 2005; Volkov *et al.* 2005). In addition, conclusions on considerable food competition between two most abundant Pacific salmon species (chum and pink salmon) are not supported by our data (Dulepova 1998; Dulepova and Dulepov *in press*; Efimkin *et al.* 2004; Temnykh, *et al.* 2004; Shuntov and Temnykh 2005).

REFERENCES

- Bugaev, A. 2003. To the Problem about Accuracy using Scale Pattern Analysis for Identification of some Local Stocks of Sockeye Salmon *Oncorhynchus nerka* from Asia and North America. (NPAFC Doc. 699). 26p. KamchatNIRO, Kamchatka Fisheries and Oceanography Research Inst., Fisheries State Commit. of Russia, Petropavlovsk-Kamchatsky, 18, Naberezhnaya Street, Petropavlovsk-Kamchatsky, Russia.
- Bugaev, A. 2004a. Scale Pattern Analysis of Age Composition of Sockeye Salmon *Oncorhynchus nerka* in R/V TINRO Trawl Catches in the Western Bering Sea in September–October 2002. (NPAFC Doc. 763). 26p. KamchatNIRO, Kamchatka Fisheries & Oceanography Research Inst., Fisheries State Commit. of Russia, 18, Naberezhnaya Street, Petropavlovsk-Kamchatsky, Russia.
- Bugaev, A. 2004b. Scale Pattern Analysis Estimates of Age and Stock Composition of Chinook Salmon *Oncorhynchus tshawytscha* in R/V TINRO Trawl Catches in Western Bering Sea in September–October 2002. (NPAFC Doc. 764). 15p. KamchatNIRO, Kamchatka Fisheries & Oceanography Research Inst., Fisheries State Commit. of Russia, 18, Naberezhnaya Street, Petropavlovsk-Kamchatsky, Russia.
- Dulepova, E.P. 1998. Trophic relationships between abundant species of salmon in the Bering Sea in summer. *Izv. TINRO* 124: 614–623. (In Russian)
- Dulepova, E.P., and V.I. Dulepov. *In press*. Trophic status of salmon in the Far Eastern seas. Annual and regional analysis of chum salmon *Oncorhynchus keta* feeding in the Bering Sea and adjacent Pacific Ocean areas. *Vopr. Ichthyologii*. (In Russian)
- Efimkin, A.Ya., A.F. Volkov, and N.A. Kuznetsova. 2004. Feeding of Pacific salmon in the Bering Sea in autumn 2003. *Izv. TINRO* 139. P. 370–387.
- Khen, G.V., and E.O. Basyuk. 2005. Oceanographic condition of the Bering Sea in BASIS. *N. Pac. Anadr. Fish Comm. Tech. Rep.* 6: 21–23.
- Shuntov, V.P., and O.S. Temnykh. 2005. North Pacific Ocean carrying capacity—Is it really too low for highly abundant salmon stocks? Myths and reality. *N. Pac. Anadr. Fish Comm. Tech. Rep.* 6: 3–7.
- Sviridov, V., I. Glebov, and V. Kulik. 2004a. Spatio-temporal variability in biological characteristics of Pacific salmon in the western Bering Sea. (NPAFC Doc. 753). 17p. Pacific Scientific Research Fisheries Centre (TINRO-Centre), 4, Shevchenko Alley, Vladivostok, 690950, Russia
- Sviridov, V., I. Glebov, M. Ocheretyanny, and V. Kulik. 2004b. Traumatization and infestation of the Pacific salmon in the western Bering Sea and adjacent Pacific waters during summer-autumn period of 2003. (NPAFC Doc. 752 Rev. 1). 17p. Pacific Scientific Research Fisheries Centre (TINRO-Centre), 4, Shevchenko Alley, Vladivostok, 690950, Russia
- Temnykh O.S. 2004. Asian pink salmon during the marine period of life: biology, spatial differentiation, role and place in pelagic communities. Doctor of Sciences Thesis. Vladivostok. TINRO-Centre. 435 pp.
- Temnykh O.S., Glebov I.I., Naidenko S.V., Starovoytov A.N., Efimkin A.Ya., Sviridov V.V., Rassadnikov O.A., Kuznetsova N.A. 2004. Contemporary status of Pacific salmon in the pelagic communities of the Far Eastern Seas. *Izv. TINRO*. Vol. 137. P. 28–44.
- Temnykh, O., I. Glebov, A. Starovoytov, and V. Sviridov. 2004. Russia Proposed Cruise Plan of R/V TINRO for Pacific Salmon Marine Period of Life Research in the Bering Sea in June–October 2004. (NPAFC Doc. 754). 5p. Laboratory of Applied biocenology, Pacific Scientific Research Fisheries Centre (TINRO-Centre), 4, Shevchenko Alley, Vladivostok, 690950, Russia.
- Volkov, A.F., A.J. Efimkin, N.A. Kuznetsova, and A.M. Slabinsky. 2005. Hydrobiological Investigations by the TINRO-Centre under the BASIS-2003 Program: zooplankton and Pacific salmon feeding. *N. Pac. Anadr. Fish Comm. Tech. Rep.* 6: 35–37.